Warren D. King, M.D.

Orthopedic Surgeon Specializing in Arthroscopy and Sports Medicine

Medial Epicondylitis (Golfer's Elbow)

What is it? Medial epicondylitis occurs with pain and inflammation on the inside of the elbow where the muscles that flex the wrist attach to the bone. This can occur not only in golfers but in anyone who performs repeated motions of the wrist. See Figure 1.

What are the symptoms?

Commonly this occurs with pain and tenderness on the inside or medial side of the elbow, pain or weakness with gripping or twisting of the wrist, and pain with lifting objects.

What causes this? Epicondylitis is caused by repetitive stress and strain to the muscles and tendons that attach the forearm muscles to the elbow. This can include any sudden change in activity level or intensity, or any incorrect grip or grip size in racquet sports. This repetitive stress causes microscopic tearing of the tendon. Blood supply to this area is poor and thus commonly the healing response is limited.

What is the treatment?

Step 1: Initially treatment consists of anti-inflammatory medication and a stretching and strengthening program. **Stretching is a key component in treating epicondylitis. Each stretch should be held for at least 10 seconds, then relax the arm and repeat up to several times a day and in between activities. Minimally 3 sets of 20 repetitions should be performed.** See Figure 2. Strengthening exercises will also help, see Figures 3-4. While doing any strengthening, exercises should always be pain free and done with a light weight and at least 3 sets of 20 repetitions.



Figure 1: From Jobe FW: Operative Techniques in Upper Extremity Sports Injuries. St. Louis, Mosby Year Book, 1996, p.441.



Figure 2: Stretching

Affiliations

Oakland Raiders Director of Orthopedic Surgery

San Francisco Giants Orthopedic Surgical Consultant

San Jose Sharks Asst. Director of Orthopedic Surgery

US Soccer National Teams Programs

US Rugby Director of Orthopedic Surgery

Palo Alto Medical Foundation

Palo Alto Office 795 El Camino Real Palo Alto, CA 94301 ph (650) 853.2943 fax (650) 853.6094 **Step 2:** If symptoms continue after a trial of stretching and strengthening, patients may benefit from a cortisone injection. Cortisone acts by decreasing inflammation and increasing blood flow to the area which aids in healing. Most patients feel relief after one injection, but up to three cortisone injections over a period of time may be given.

Step 3: If stretching and strengthening, anti- inflammatory medication, and cortisone injections do not relieve symptoms that surgery is an option. Through a small incision over the bony prominence on the inside of the elbow the base of the diseased tendon is removed and then normal tendon tissue is used to reattach to the bone. This is an outpatient surgery.



Figure 4: Supination and Pronation. 3 sets of 20 repetitions 3-4 days a week, pain free.



Figure 3: Wrist Flexors. 3 sets of 20 repetitions 3-4 days a week, pain free.